

---

## Exercises

**4.3 Invariant measures.**<sup>1,2</sup> (Mathematics, Complexity) ④

Liouville's theorem tells us that all available points in phase space are equally weighted when a Hamiltonian system is averaged over all times. What happens for systems that evolve according to laws that are not Hamiltonian? Usually, the system does not continue to explore all points in its state space; at long times it is confined to a subset of the original space known as the *attractor*.

We consider the behavior of the 'logistic' mapping from the unit interval  $(0, 1)$  into itself:<sup>3</sup>

$$f(x) = 4\mu x(1 - x). \quad (4.12)$$

We talk of the trajectory of an initial point  $x_0$  as the sequence of points  $x_0, f(x_0), f(f(x_0)), \dots, f^{[n]}(x_0), \dots$ . Iteration can be thought of as a time step (one iteration of a Poincaré return map of Exercise 4.4 or one step  $\Delta t$  in a time-step algorithm as in Exercise 3.12).

*Attracting fixed point.* For small  $\mu$ , our mapping has an attracting fixed-point. A fixed-point of a mapping is a value  $x^* = f(x^*)$ ; a fixed-point is stable if small perturbations shrink after iterating:

$$|f(x^* + \epsilon) - x^*| \approx |f'(x^*)|\epsilon < \epsilon, \quad (1)$$

which happens if the derivative  $|f'(x^*)| < 1$ .<sup>4</sup>

(a) Iteration. Set  $\mu = 0.2$ ; iterate  $f$  for some initial points  $0 < x_0 < 1$  of your choosing, and convince yourself that they are all attracted to zero. Plot  $f$  and the diagonal  $y = x$  on the same plot. Are there any fixed-points other than  $x = 0$ ? Repeat for  $\mu = 0.4$ , and 0.6. What happens?

Analytics. Find the non-zero fixed-point  $x^*(\mu)$  of the map 4.12, and show that it exists and is stable for  $1/4 < \mu < 3/4$ . If you are ambitious or have a computer algebra program, show that there is a stable, period-two cycle for  $3/4 < \mu < (1 + \sqrt{6})/4$ .

An attracting fixed-point is the antithesis of Liouville's theorem; all initial conditions are transient except one, and all systems lead eventually to the same, time-independent state. (On the other hand, this is precisely the behavior we expect in statistical mechanics *on the macroscopic scale*; the system settles down into a time-independent equilibrium state! All microstates are equivalent, but the vast majority of accessible microstates have the same macroscopic behavior in most large systems.) We could define a rather trivial 'equilibrium ensemble' for this system, which consists of the single point  $x^*$ ; any property  $O(x)$  will have the long-time average  $\langle O \rangle = O(x^*)$ .

For larger values of  $\mu$ , more complicated things happen. At  $\mu = 1$ , the dynamics can be shown to fill the entire interval; the dynamics is ergodic, and the attractor fills the entire set of available states. However, unlike the case of Hamiltonian systems, not all states are weighted equally (i.e., Liouville's theorem does not hold).

We can find time averages for functions of  $x$  in two ways: by averaging over time (many iterates of the map) or by weighting an integral over  $x$  by the *invariant density*  $\rho(x)$ . The invariant density  $\rho(x) dx$  is the probability that a point on a long trajectory will lie between  $x$  and  $x + dx$ . To find it numerically, we iterate a typical point<sup>5</sup>  $x_0$  a thousand

<sup>1</sup>From *Statistical Mechanics: Entropy, Order Parameters, and Complexity* by James P. Sethna, copyright Oxford University Press, 2007, page 70. A pdf of the text is available at [pages.physics.cornell.edu/sethna/StatMech/](http://pages.physics.cornell.edu/sethna/StatMech/) (select the picture of the text). Hyperlinks from this exercise into the text will work if the latter PDF is downloaded into the same directory/folder as this PDF.

<sup>2</sup>This exercise and the associated software were developed in collaboration with Christopher Myers; see [64]. Hints available [129].

<sup>3</sup>We also study this map in Exercises 5.9, 5.16, and 12.9.

<sup>4</sup>For many-dimensional mappings, a sufficient criterion for stability is that all the eigenvalues of the Jacobian have magnitude smaller than one. A continuous time evolution  $dy/dt = F(y)$  will be stable if  $dF/dy$  is smaller than zero, or (for multidimensional systems) if the Jacobian  $DF$  has eigenvalues whose real parts are all less than zero.

<sup>5</sup>For example, we must not choose an unstable fixed-point or unstable periodic orbit!

or so times ( $N_{\text{transient}}$ ) to find a point  $x_a$  on the attractor, and then collect a long trajectory of perhaps a million points ( $N_{\text{cycles}}$ ). A histogram of this trajectory gives  $\rho(x)$ . Averaging over this density is manifestly the same as a time average over the trajectory of a million points. We call  $\rho(x)$  invariant because it is left the same under the mapping  $f$ ; iterating our million-point approximation for  $\rho$  once under  $f$  only removes the first point  $x_a$  and adds one extra point to the end.

(b) Invariant density. Set  $\mu = 1$ ; iterate  $f$  many times, and form a histogram of values giving the density  $\rho(x)$  of points along the trajectory. You should find that points  $x$  near the boundaries are approached more often than points near the center.

*Analytics.* Using the fact that the long-time average  $\rho(x)$  must be independent of time, verify for  $\mu = 1$  that the density of points is<sup>6</sup>

$$\rho(x) = \frac{1}{\pi\sqrt{x(1-x)}}. \quad (4.14)$$

Plot this theoretical curve with your numerical histogram. (Hint: The points in a range  $dx$  around a point  $x$  map under  $f$  to a range  $dy = f'(x)dx$  around the image  $y = f(x)$ . Each iteration maps two points  $x_a$  and  $x_b = 1 - x_a$  to  $y$ , and thus maps all the density  $\rho(x_a)|dx_a|$  and  $\rho(x_b)|dx_b|$  into  $dy$ . Hence the probability  $\rho(y)dy$  must equal  $\rho(x_a)|dx_a| + \rho(x_b)|dx_b|$ , so

$$\rho(f(x_a)) = \rho(x_a)/|f'(x_a)| + \rho(x_b)/|f'(x_b)|. \quad (4.15)$$

Substitute eqn 4.14 into eqn 4.15. You will need to factor a polynomial.)


Mathematicians call this probability density  $\rho(x)dx$  the *invariant measure* on the attractor.<sup>7</sup> To get the long-term average of any function  $O(x)$ , one can use

$$\langle O \rangle = \int O(x)\rho(x)dx. \quad (2)$$

To a mathematician, a measure is a way of weighting different regions when calculating integrals—precisely our  $\rho(x)dx$ . Notice that, for the case of an attracting fixed-point, we would have  $\rho(x) = \delta(x - x^*)$ .<sup>8</sup>

*Cusps in the invariant density.* At values of  $\mu$  slightly smaller than one, our mapping has a rather complex invariant density.


(c) Find the invariant density (as described above) for  $\mu = 0.9$ . Make your trajectory length  $N_{\text{cycles}}$  big enough and the bin size small enough to see the interesting structures. Notice that the attractor no longer fills the whole range  $(0, 1)$ ; locate roughly where the edges are. Notice also the cusps in  $\rho(x)$  at the edges of the attractor, and also at places inside the attractor (called boundaries, see [64]). Locate some of the more prominent cusps.



**Fig. 4.4** Invariant density in the chaotic region ( $\mu = 0.95$ ).

*Analytics of cusps.* Notice that  $f'(1/2) = 0$ , so by eqn 4.15 we know that  $\rho(f(x)) \geq \rho(x)/|f'(x)|$  must have a singularity near  $x = 1/2$ ; all the points near  $x = 1/2$  are squeezed together and folded to one side by  $f$ . Further iterates of this singularity produce more cusps; the crease after one fold stays a crease after being further stretched and kneaded.

(d) Set  $\mu = 0.9$ . Calculate  $f(1/2)$ ,  $f(f(1/2))$ , ... and compare these iterates to the locations of the edges and cusps from part (c). (You may wish to include them both on the same plot.)



<sup>6</sup>You need not derive the factor  $1/\pi$ , which normalizes the probability density to one.

<sup>7</sup>There are actually many possible invariant measures on some attractors; this one is the SRB measure (John Guckenheimer, private communication).

<sup>8</sup>The case of a fixed-point then becomes mathematically a measure with a point mass at  $x^*$ .

**Fig. 4.5 Bifurcation diagram in the chaotic region.** Notice the boundary lines threading through the diagram, images of the crease formed by the folding at  $x = \frac{1}{2}$  in our map (see [64]).

*Bifurcation diagram.* The evolution of the attractor and its invariant density as  $\mu$  varies are plotted in the *bifurcation diagram*, which is shown for large  $\mu$  in Fig. 4.5. One of the striking features in this plot are the sharp boundaries formed by the cusps.

(e) Bifurcation diagram. Plot the attractor (duplicating Fig. 4.5) as a function of  $\mu$ , for  $0.8 < \mu < 1$ . (Pick regularly-spaced  $\delta\mu$ , run  $n_{\text{transient}}$  steps, record  $n_{\text{cycles}}$  steps, and plot. After the routine is working, you should be able to push  $n_{\text{transient}}$  and  $n_{\text{cycles}}$  both larger than 100, and  $\delta\mu < 0.01$ .)

On the same plot, for the same  $\mu$ s, plot the first eight images of  $x = \frac{1}{2}$ , that is,  $f(\frac{1}{2}), f(f(\frac{1}{2})), \dots$ . Are the boundaries you see just the cusps? What happens in the bifurcation diagram when two boundaries touch? (See [64].)